

SOLUTIONS WITH INTERSECTING P-BRANES RELATED TO TODA CHAINS

V.D. Ivashchuk¹, and S.-W. Kim ²

*Center for Gravitation and Fundamental Metrology, VNIIMS, 3/1 M. Ulyanovoy Str.,
Moscow 117313, Russia¹*

*Department of Science Education and Basic Science Research Institute, Ewha Womans
University, Seoul 120-750, Korea²*

Abstract

Solutions in multidimensional gravity with m p -branes related to Toda-like systems (of general type) are obtained. These solutions are defined on a product of $n + 1$ Ricci-flat spaces $M_0 \times M_1 \times \dots \times M_n$ and are governed by one harmonic function on M_0 . The solutions are defined up to the solutions of Laplace and Toda-type equations and correspond to null-geodesics of the (sigma-model) target-space metric. Special solutions relating to A_m Toda chains (e.g. with $m = 1, 2$) are considered.

¹ivas@rgs.phys.msu.su

²sungwon@mm.ewha.ac.kr

I Introduction

At present there exists a special interest to the so-called M theory (see, for example, [1]-[2]). This theory is “supermembrane” analogue of superstring models [3] in $D = 11$. The low-energy limit of M -theory after a dimensional reduction leads to models governed by a Lagrangian containing metric, fields of forms and scalar fields. These models contain a large variety of so-called p -brane solutions (see [4]-[42] and references therein).

In [25] it was shown that after dimensional reduction on the manifold $M_0 \times M_1 \times \dots \times M_n$ and when the composite p -brane ansatz for fields of forms is considered the problem is reduced to the gravitating self-interacting σ -model with certain constraints imposed. (For electric p -branes see also [16, 17, 26].) This representation may be considered as a powerful tool for obtaining different solutions with intersecting p -branes (analogues of membranes). In [25, 26, 36, 37, 38] the Majumdar-Papapetrou type solutions (see [45]) were obtained (for non-composite case see [16, 17]). These solutions correspond to Ricci-flat (M_i, g^i) , (g^i is metric on M_i) $i = 1, \dots, n$, and were also generalized to the case of Einstein internal spaces [25]. Earlier some special classes of these solutions were considered in [8, 9, 11, 19, 20, 21]. The obtained solutions take place, when certain (block-)orthogonality relations (on couplings parameters, dimensions of “branes”, total dimension) are imposed. In this situation a class of cosmological and spherically-symmetric solutions was obtained [31, 40]. Special cases were also considered in [12, 27, 29, 30]. The solutions with the horizon were considered in details in [7, 22, 23, 31, 33].

In models under consideration there exists a large variety of Toda-chain solutions, when certain intersection rules are satisfied [31]. Cosmological and spherically symmetric solutions with p -branes and n internal spaces related to A_m Toda chains were previously considered in [12, 13] and [39, 42].

It is well known that geodesics of the target space equipped with some harmonic function on a three-dimensional space generate a solution to the σ -model equations [46, 47]. (It was observed in [48] that null geodesics of the target space of stationary five-dimensional Kaluza-Klein theory may be used to generate multisoliton solutions similar to the Israel-Wilson-Perjès solutions of Einstein-Maxwell theory.) Here we apply this null-geodesic method to our sigma-model and obtain a new class of solutions in multidimensional gravity with p -branes governed by one harmonic function H . The solutions from this class correspond to null-geodesics of the target-space metric and are defined by some functions $f_s(H) = \exp(-q^s(H))$ with $q^s(u)$ being solutions to Toda-type equations.

II The model

We consider a model governed by the action [25]

$$S = \int d^D z \sqrt{|g|} \left\{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \sum_{a \in \Delta} \frac{\theta_a}{n_a!} \exp[2\lambda_a(\varphi)] (F^a)^2 \right\} \quad (2.1)$$

where $g = g_{MN} dz^M \otimes dz^N$ is a metric, $\varphi = (\varphi^\alpha) \in \mathbf{R}^l$ is a vector of scalar fields, $(h_{\alpha\beta})$ is a constant symmetric non-degenerate $l \times l$ matrix ($l \in \mathbf{N}$), $\theta_a = \pm 1$, $F^a = dA^a$ is a

n_a -form ($n_a \geq 1$), λ_a is a 1-form on $\mathbf{R}^l$: $\lambda_a(\varphi) = \lambda_{\alpha a} \varphi^\alpha$, $a \in \Delta$, $\alpha = 1, \dots, l$. Here Δ is some finite set.

We consider a manifold

$$M = M_0 \times M_1 \times \dots \times M_n, \quad (2.2)$$

with a metric

$$g = e^{2\gamma(x)} g^0 + \sum_{i=1}^n e^{2\phi^i(x)} g^i \quad (2.3)$$

where $g^0 = g_{\mu\nu}^0(x) dx^\mu \otimes dx^\nu$ is a metric on the manifold M_0 , and $g^i = g_{m_i n_i}^i(y_i) dy_i^{m_i} \otimes dy_i^{n_i}$ is an Einstein metric on M_i satisfying the equation

$$R_{m_i n_i}[g^i] = \xi_i g_{m_i n_i}^i, \quad (2.4)$$

$m_i, n_i = 1, \dots, d_i$; $\xi_i = \text{const}$, $i = 1, \dots, n$. (Here we identify notations for g^i and $\hat{g}^i$, where $\hat{g}^i = p_i^* g^i$ is the pullback of the metric g^i to the manifold M by the canonical projection: $p_i : M \rightarrow M_i$, $i = 0, \dots, n$. An analogous agreement will be also kept for volume forms etc.)

Any manifold M_ν is supposed to be oriented and connected and $d_\nu \equiv \dim M_\nu$, $\nu = 0, \dots, n$. Let

$$\tau_i \equiv \sqrt{|g^i(y_i)|} dy_i^1 \wedge \dots \wedge dy_i^{d_i}, \quad \varepsilon(i) \equiv \text{sign}(\det(g_{m_i n_i}^i)) = \pm 1 \quad (2.5)$$

denote the volume d_i -form and signature parameter respectively, $i = 1, \dots, n$. Let $\Omega = \Omega_n$ be a set of all subsets of $\{1, \dots, n\}$, $|\Omega| = 2^n$. For any $I = \{i_1, \dots, i_k\} \in \Omega$, $i_1 < \dots < i_k$, we denote

$$\tau(I) \equiv \tau_{i_1} \wedge \dots \wedge \tau_{i_k}, \quad d(I) \equiv \sum_{i \in I} d_i, \quad \varepsilon(I) \equiv \prod_{i \in I} \varepsilon(i). \quad (2.6)$$

We also put $\tau(\emptyset) = \varepsilon(\emptyset) = 1$ and $d(\emptyset) = 0$.

For fields of forms we consider the following composite electromagnetic ansatz

$$F^a = \sum_{I \in \Omega_{a,e}} \mathcal{F}^{(a,e,I)} + \sum_{J \in \Omega_{a,m}} \mathcal{F}^{(a,m,J)} \quad (2.7)$$

where

$$\mathcal{F}^{(a,e,I)} = d\Phi^{(a,e,I)} \wedge \tau(I), \quad (2.8)$$

$$\mathcal{F}^{(a,m,J)} = e^{-2\lambda_a(\varphi)} * (d\Phi^{(a,m,J)} \wedge \tau(J)) \quad (2.9)$$

are elementary forms of electric and magnetic types respectively, $a \in \Delta$, $I \in \Omega_{a,e}$, $J \in \Omega_{a,m}$ and $\Omega_{a,e}$, $\Omega_{a,m}$ are non-empty subsets of Ω . In (2.9) $*$ = $*[g]$ is the Hodge operator on (M, g) . For scalar functions we put

$$\varphi^\alpha = \varphi^\alpha(x), \quad \Phi^s = \Phi^s(x), \quad (2.10)$$

$s \in S$, $x \in M_0$.

Here and below

$$S = S_e \cup S_m, \quad S_v = \bigcup_{a \in \Delta} \{a\} \times \{v\} \times \Omega_{a,v}, \quad (2.11)$$

$v = e, m$. The set S consists of elements $s = (a_s, v_s, I_s)$, where $a_s \in \Delta$, $v_s = e, m$ and $I_s \in \Omega_{a_s, v_s}$.

Due to (2.8) and (2.9)

$$d(I) = n_a - 1, \quad d(J) = D - n_a - 1, \quad (2.12)$$

for $I \in \Omega_{a,e}$, $J \in \Omega_{a,m}$.

II.1 The sigma model

Let $d_0 \neq 2$ and

$$\gamma = \gamma_0(\phi) \equiv \frac{1}{2 - d_0} \sum_{j=1}^n d_j \phi^j, \quad (2.13)$$

i.e. the generalized harmonic gauge is used.

We impose the restriction on sets $\Omega_{a,v}$. These restrictions guarantee the block-diagonal structure of a stress-energy tensor (like for the metric) and the existence of σ -model representation [25].

We denote $w_1 \equiv \{i | i \in \{1, \dots, n\}, d_i = 1\}$, and $n_1 = |w_1|$ (i.e. n_1 is the number of 1-dimensional spaces among M_i , $i = 1, \dots, n$).

Restriction 1. Let 1a) $n_1 \leq 1$ or 1b) $n_1 \geq 2$ and for any $a \in \Delta$, $v \in \{e, m\}$, $i, j \in w_1$, $i < j$, there are no $I, J \in \Omega_{a,v}$ such that $i \in I$, $j \in J$ and $I \setminus \{i\} = J \setminus \{j\}$.

Restriction 2 (only for $d_0 = 1, 3$). Let 2a) $n_1 = 0$ or 2b) $n_1 \geq 1$ and for any $a \in \Delta$, $i \in w_1$ there are no $I \in \Omega_{a,m}$, $J \in \Omega_{a,e}$ such that $\bar{I} = \{i\} \sqcup J$ for $d_0 = 1$ and $J = \{i\} \sqcup \bar{I}$ for $d_0 = 3$. Here and in what follows

$$\bar{I} \equiv \{1, \dots, n\} \setminus I. \quad (2.14)$$

These restrictions are satisfied in the non-composite case [16, 17]: $|\Omega_{a,v}| = 1$, (i.e. when there are no two p -branes with the same color index a , $a \in \Delta$.) Restriction 1 and 2 forbid certain intersections of two p -branes with the same color index for $n_1 \geq 2$ and $n_1 \geq 1$ respectively.

It was proved in [25] that equations of motion for the model (2.1) and the Bianchi identities: $d\mathcal{F}^s = 0$, $s \in S_m$, for fields from (2.3)–(2.13), when Restrictions 1 and 2 are imposed, are equivalent to equations of motion for the σ -model governed by the action

$$S_\sigma = \int d^{d_0}x \sqrt{|g^0|} \{ R[g^0] - \hat{G}_{AB} g^{0\mu\nu} \partial_\mu z^A \partial_\nu z^B - \sum_{s \in S} \varepsilon_s e^{-2U_A^s z^A} g^{0\mu\nu} \partial_\mu \Phi^s \partial_\nu \Phi^s - 2V \}, \quad (2.15)$$

where $(z^A) = (\phi^i, \varphi^\alpha)$, the index set S is defined in (2.11),

$$V = V(\phi) = -\frac{1}{2} \sum_{i=1}^n \xi_i d_i e^{-2\phi^i + 2\gamma_0(\phi)} \quad (2.16)$$

is the potential,

$$(\hat{G}_{AB}) = \begin{pmatrix} G_{ij} & 0 \\ 0 & h_{\alpha\beta} \end{pmatrix}, \quad (2.17)$$

is the target space metric with

$$G_{ij} = d_i \delta_{ij} + \frac{d_i d_j}{d_0 - 2}, \quad (2.18)$$

$$(U_A^s) = (d_i \delta_{iI_s}, -\chi_s \lambda_{\alpha a_s}), \quad (2.19)$$

are vectors, $s = (a_s, v_s, I_s)$, $\chi_e = +1$, $\chi_m = -1$;

$$\delta_{iI} = \sum_{j \in I} \delta_{ij} \quad (2.20)$$

is the indicator of i belonging to I : $\delta_{iI} = 1$ for $i \in I$ and $\delta_{iI} = 0$ otherwise; and

$$\varepsilon_s = (-\varepsilon[g])^{(1-\chi_s)/2} \varepsilon(I_s) \theta_{a_s}, \quad (2.21)$$

$s \in S$, $\varepsilon[g] \equiv \text{sign det}(g_{MN})$. More explicitly (2.21) reads $\varepsilon_s = \varepsilon(I_s) \theta_{a_s}$ for $v_s = e$ and $\varepsilon_s = -\varepsilon[g] \varepsilon(I_s) \theta_{a_s}$, for $v_s = m$.

III Exact solutions with one harmonic function

III.1 Toda-like Lagrangian

Action (2.15) may be also written in the form

$$S_\sigma = \int d^{d_0} x \sqrt{|g^0|} \{ R[g^0] - \mathcal{G}_{\hat{A}\hat{B}}(X) g^{0\mu\nu} \partial_\mu X^{\hat{A}} \partial_\nu X^{\hat{B}} - 2V \} \quad (3.1)$$

where $X = (X^{\hat{A}}) = (\phi^i, \varphi^\alpha, \Phi^s) \in \mathbf{R}^N$, and minisupermetric

$$\mathcal{G} = \mathcal{G}_{\hat{A}\hat{B}}(X) dX^{\hat{A}} \otimes dX^{\hat{B}} \quad (3.2)$$

on minisuperspace

$$\mathcal{M} = \mathbf{R}^N, \quad N = n + l + |S| \quad (3.3)$$

($|S|$ is the number of elements in S) is defined by the relation

$$(\mathcal{G}_{\hat{A}\hat{B}}(X)) = \begin{pmatrix} G_{ij} & 0 & 0 \\ 0 & h_{\alpha\beta} & 0 \\ 0 & 0 & \varepsilon_s \exp(-2U^s(X)) \delta_{ss'} \end{pmatrix}. \quad (3.4)$$

Here we consider exact solutions to field equations corresponding to the action (3.1)

$$R_{\mu\nu}[g^0] = \mathcal{G}_{\hat{A}\hat{B}}(X) \partial_\mu X^{\hat{A}} \partial_\nu X^{\hat{B}} + \frac{2V}{d_0 - 2} g_{\mu\nu}^0, \quad (3.5)$$

$$\frac{1}{\sqrt{|g^0|}} \partial_\mu [\sqrt{|g^0|} \mathcal{G}_{\hat{C}\hat{B}}(X) g^{0\mu\nu} \partial_\nu X^{\hat{B}}] - \frac{1}{2} \mathcal{G}_{\hat{A}\hat{B},\hat{C}}(X) g^{0,\mu\nu} \partial_\mu X^{\hat{A}} \partial_\nu X^{\hat{B}} = V_{,\hat{C}}, \quad (3.6)$$

$s \in S$. Here $\Delta[g^0]$ is the Laplace-Beltrami operator corresponding to g^0 and $V_{,\hat{C}} = \partial V / \partial X^{\hat{C}}$.

We put

$$X^{\hat{A}}(x) = F^{\hat{A}}(H(x)), \quad (3.7)$$

where $F : (u_-, u_+) \rightarrow \mathbf{R}^N$ is a smooth function, $H : M_0 \rightarrow \mathbf{R}$ is a harmonic function on M_0 , i.e.

$$\Delta[g^0]H = 0, \quad (3.8)$$

satisfying $u_- < H(x) < u_+$ for all $x \in M_0$.

The substitution of (3.7) into eqs. (3.5) and (3.6) leads us to the relations

$$R_{\mu\nu}[g^0] = \mathcal{G}_{\hat{A}\hat{B}}(F(u)) \dot{F}^{\hat{A}} \dot{F}^{\hat{B}} \partial_\mu H \partial_\nu H + \frac{2V}{d_0 - 2} g_{\mu\nu}^0, \quad (3.9)$$

$$\left[\frac{d}{du} (\mathcal{G}_{\hat{C}\hat{B}}(F(u)) \dot{F}^{\hat{B}}) - \frac{1}{2} \mathcal{G}_{\hat{A}\hat{B},\hat{C}}(F(u)) \dot{F}^{\hat{A}} \dot{F}^{\hat{B}} \right] g^{0,\mu\nu} \partial_\mu H \partial_\nu H = V_{,\hat{C}}, \quad (3.10)$$

where $u = H(x)$ and $\dot{f} = df/du$.

Let all spaces (M_i, g^i) be Ricci-flat, i.e.

$$R_{m_i n_i}[g^i] = 0, \quad (3.11)$$

$i = 0, \dots, n$. In this case the potential is zero : $V = 0$ and the field equations (3.9) and (3.10) are satisfied identically if $F = F(u)$ obey the Lagrange equations for the Lagrangian

$$L = \frac{1}{2} \mathcal{G}_{\hat{A}\hat{B}}(F) \dot{F}^{\hat{A}} \dot{F}^{\hat{B}} \quad (3.12)$$

with the zero-energy constraint

$$E = \frac{1}{2} \mathcal{G}_{\hat{A}\hat{B}}(F) \dot{F}^{\hat{A}} \dot{F}^{\hat{B}} = 0. \quad (3.13)$$

This means that $F : (u_-, u_+) \rightarrow \mathbf{R}^N$ is a null-geodesic map for the minisupermetric (3.2). Thus, we are led to the Lagrange system (3.12) with the minisupermetric $\mathcal{G}$ defined in (3.4).

The problem of integrability may be simplified if we integrate the Lagrange equations corresponding to Φ^s (i.e. the Maxwell equations for $s \in S_e$ and Bianchi identities for $s \in S_m$):

$$\frac{d}{du} (\exp(-2U^s(z)) \dot{\Phi}^s) = 0 \iff \dot{\Phi}^s = Q_s \exp(2U^s(z)), \quad (3.14)$$

where Q_s are constants, $s \in S$. Here $(F^{\hat{A}}) = (z^A, \Phi^s)$. We put

$$Q_s \neq 0, \quad (3.15)$$

for all $s \in S$.

For fixed $Q = (Q_s, s \in S)$ the Lagrange equations for the Lagrangian (3.12) corresponding to $(z^A) = (\phi^i, \varphi^\alpha)$, when equations (3.14) are substituted, are equivalent to the Lagrange equations for the Lagrangian

$$L_Q = \frac{1}{2} \hat{G}_{AB} \dot{z}^A \dot{z}^B - V_Q, \quad (3.16)$$

where

$$V_Q = \frac{1}{2} \sum_{s \in S} \varepsilon_s Q_s^2 \exp[2U^s(z)], \quad (3.17)$$

$(\hat{G}_{AB})$ are defined in (2.17) respectively. The zero-energy constraint (3.13) reads

$$E_Q = \frac{1}{2} \hat{G}_{AB} \dot{z}^A \dot{z}^B + V_Q = 0. \quad (3.18)$$

III.2 Toda-type solutions

Let us define the scalar product as follows

$$(U, U') = \hat{G}^{AB} U_A U'_B, \quad (3.19)$$

for $U, U' \in \mathbf{R}^{n+l}$, where $(\hat{G}^{AB}) = (\hat{G}_{AB})^{-1}$. The scalar products (3.19) for vectors U^s were calculated in [25]

$$(U^s, U^{s'}) = d(I_s \cap I_{s'}) + \frac{d(I_s)d(I_{s'})}{2-D} + \chi_s \chi_{s'} \lambda_{\alpha a_s} \lambda_{\beta a_{s'}} h^{\alpha\beta}, \quad (3.20)$$

where $(h^{\alpha\beta}) = (h_{\alpha\beta})^{-1}$; $s = (a_s, v_s, I_s)$ and $s' = (a_{s'}, v_{s'}, I_{s'})$ belong to S .

Here we are interested in exact solutions for a special case when the vectors $U^s \in \mathbf{R}^{n+l}$ satisfy the following conditions

$$K_s = (U^s, U^s) \neq 0, \quad (3.21)$$

for all $s \in S$, and a (“quasi-Cartan”) matrix

$$(A_{ss'}) = \left(\frac{2(U^s, U^{s'})}{(U^{s'}, U^{s'})} \right) \quad (3.22)$$

is a non-degenerate one. Here some ordering in S is assumed. It follows from (3.21) and the non-degeneracy of the matrix (3.22) that the vectors $U^s, s \in S$, are linearly independent. Hence, the number of the vectors U^s should not exceed the dimension of $\mathbf{R}^{n+l}$, i.e.

$$|S| \leq n + l. \quad (3.23)$$

From (3.20)-(3.22) we get the following intersection rules [31]

$$d(I_s \cap I_{s'}) = \frac{d(I_s)d(I_{s'})}{D-2} - \chi_s \chi_{s'} \lambda_{a_s} \cdot \lambda_{a_{s'}} + \frac{1}{2} K_{s'} A_{ss'}, \quad (3.24)$$

$s \neq s'$.

The exact solutions to Lagrange equations corresponding to (3.16) with the potential (3.17) could be readily obtained using the relations from Appendix. The solutions read

$$z^A = \sum_{s \in S} \frac{U^{sA}}{(U^s, U^s)} q^s + c^A u + \bar{c}^A, \quad (3.25)$$

where q^s are solutions to Toda-type equations

$$\ddot{q}^s = -B_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right), \quad (3.26)$$

with

$$B_s = 2K_s A_s, \quad A_s = \frac{1}{2} \varepsilon_s Q_s^2, \quad (3.27)$$

$s \in S$. These equations correspond to the Lagrangian

$$L_{TL} = \frac{1}{4} \sum_{s, s' \in S} K_s^{-1} A_{ss'} \dot{q}^s \dot{q}^{s'} - \sum_{s \in S} A_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right). \quad (3.28)$$

Vectors $c = (c^A)$ and $\bar{c} = (\bar{c}^A)$ satisfy the linear constraint relations (see (6.11) in Appendix)

$$U^s(c) = U_A^s c^A = \sum_{i \in I_s} d_i c^i - \chi_s \lambda_{a_s \alpha} c^\alpha = 0, \quad (3.29)$$

$$U^s(\bar{c}) = U_A^s \bar{c}^A = \sum_{i \in I_s} d_i \bar{c}^i - \chi_s \lambda_{a_s \alpha} \bar{c}^\alpha = 0, \quad (3.30)$$

$s \in S$.

The contravariant components $U^{rA} = \hat{G}^{AB} U_B^r$ are [25, 31]

$$U^{si} = G^{ij} U_j^s = \delta_{iI_s} - \frac{d(I_s)}{D-2}, \quad U^{s\alpha} = -\chi_s \lambda_{a_s}^\alpha, \quad (3.31)$$

Here (as in [49])

$$G^{ij} = \frac{\delta^{ij}}{d_i} + \frac{1}{2-D}, \quad (3.32)$$

$i, j = 1, \dots, n$, are the components of the matrix inverse to (G_{ij}) from (2.18).

Using (3.25) and (3.31) we obtain

$$\phi^i = \sum_{s \in S} h_s \left(\delta_{iI_s} - \frac{d(I_s)}{D-2} \right) q^s + c^i u + \bar{c}^i, \quad (3.33)$$

and

$$\varphi^\alpha = - \sum_{s \in S} h_s \chi_s \lambda_{a_s}^\alpha q^s + c^\alpha u + \bar{c}^\alpha, \quad (3.34)$$

$\alpha = 1, \dots, l$, and $i = 1, \dots, n$. For γ_0 from (2.13) we get

$$\gamma_0(\phi) = - \sum_{s \in S} \frac{d(I_s)}{D-2} h_s q^s + c^0 u + \bar{c}^0, \quad (3.35)$$

where we denote

$$h_s = K_s^{-1} \quad (3.36)$$

and

$$c^0 = \frac{1}{2-d_0} \sum_{j=1}^n d_j c^j, \quad \bar{c}^0 = \frac{1}{2-d_0} \sum_{j=1}^n d_j \bar{c}^j. \quad (3.37)$$

The zero-energy constraint reads (see Appendix)

$$\begin{aligned} 2E &= 2E_{TL} + \hat{G}_{ABC}^A c^B = \\ 2E_{TL} + h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^n d_i (c^i)^2 + \frac{1}{d_0-2} \left(\sum_{i=1}^n d_i c^i \right)^2 &= 0. \end{aligned} \quad (3.38)$$

where

$$E_{TL} = \frac{1}{4} \sum_{s, s' \in S} K_s^{-1} A_{ss'} \dot{q}^s \dot{q}^{s'} + \sum_{s \in S} A_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right), \quad (3.39)$$

is an integration constant (energy) for the solutions from (3.26).

From the relation

$$\exp(2U^s(z)) = \prod_{s' \in S} f_{s'}^{-A_{ss'}}, \quad (3.40)$$

following from (3.22), (3.25), (3.29) and (3.30) we get for electric-type forms (2.8)

$$\mathcal{F}^s = Q_s \left(\prod_{s' \in S} f_{s'}^{-A_{ss'}} \right) dH \wedge \tau(I_s), \quad (3.41)$$

$s \in S_e$, and for magnetic-type forms (2.9)

$$\mathcal{F}^s = \exp[-2\lambda_a(\varphi)] * \left[Q_s \left(\prod_{s' \in S} f_{s'}^{-A_{ss'}} \right) dH \wedge \tau(I_s) \right] = \bar{Q}_s (*_0 dH) \wedge \tau(\bar{I}_s), \quad (3.42)$$

$s \in S_m$, where $\bar{Q}_s = Q_s \varepsilon(I_s) \mu(I_s)$ and $\mu(I) = \pm 1$ is defined by the relation $\mu(I) dH \wedge \tau(I_0) = \tau(\bar{I}) \wedge dH \wedge \tau(I)$, $I_0 = \{1, \dots, n\}$ (see eq. (2.26) in [25]). Here $*_0 = *[g^0]$ is the Hodge operator on (M_0, g^0) .

Relations for the metric and scalar fields follows from (3.33)-(3.35)

$$g = \left(\prod_{s \in S} f_s^{2d(I_s)h_s/(D-2)} \right) \left\{ \exp(2c^0 H + 2\bar{c}^0) g^0 \right. \quad (3.43)$$

$$\left. + \sum_{i=1}^n \left(\prod_{s \in S} f_s^{-2h_s \delta_{iI_s}} \right) \exp(2c^i H + 2\bar{c}^i) g^i \right\}, \quad (3.44)$$

$$\exp(\varphi^\alpha) = \left(\prod_{s \in S} f_s^{h_s \chi_s \lambda_{a_s}^\alpha} \right) \exp(c^\alpha H + \bar{c}^\alpha), \quad (3.45)$$

$\alpha = 1, \dots, l$. Here

$$f_s = f_s(H) = \exp(-q^s(H)), \quad (3.46)$$

where $q^s(u)$ is a solution to Toda-like equations (3.26) and $H = H(x)$ ($x \in M_0$) is a harmonic function on (M_0, g^0) (see (3.8)).

The solution is presented by relations (3.41)-(3.46) with the functions q^s defined in (3.26) and the relations on the parameters of solutions $c^A, \bar{c}^A$ ($A = i, \alpha, 0$), Q_s, h_s imposed in (3.29), (3.30), (3.37) and (3.36), respectively.

This solution describes a set of charged (by forms) overlapping p -branes ($p_s = d(I_s) - 1, s \in S$) “living” on submanifolds of $M_1 \times \dots \times M_n$. The solution is valid if the dimensions of p -branes and dilatonic coupling vector satisfy the relations (2.12).

IV Solutions corresponding to A_m Toda chain

Here we consider exact solutions to equations of motion of a Toda-chain corresponding to the Lie algebra $A_m = sl(m+1, \mathbf{C})$ [50, 51],

$$\ddot{q}^s = -B_s \exp\left(\sum_{s'=1}^m A_{ss'} q^{s'}\right), \quad (4.1)$$

where

$$(A_{ss'}) = \begin{pmatrix} 2 & -1 & 0 & \dots & 0 & 0 \\ -1 & 2 & -1 & \dots & 0 & 0 \\ 0 & -1 & 2 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 2 & -1 \\ 0 & 0 & 0 & \dots & -1 & 2 \end{pmatrix} \quad (4.2)$$

is the Cartan matrix of the Lie algebra A_m and $B_s > 0, s, s' = 1, \dots, m$. Here we put $S = \{1, \dots, m\}$.

The equations of motion (4.1) correspond to the Lagrangian

$$L_T = \frac{1}{2} \sum_{s,s'=1}^m A_{ss'} \dot{q}^s \dot{q}^{s'} - \sum_{s=1}^m B_s \exp\left(\sum_{s'=1}^m A_{ss'} q^{s'}\right). \quad (4.3)$$

This Lagrangian may be obtained from the standard one [50] by separating a coordinate describing the motion of the center of mass.

Using the result of A. Anderson [51] we present the solution to eqs. (4.1) in the following form

$$C_s \exp(-q^s(u)) = \sum_{r_1 < \dots < r_s}^{m+1} v_{r_1} \dots v_{r_s} \Delta^2(w_{r_1}, \dots, w_{r_s}) \exp[(w_{r_1} + \dots + w_{r_s})u], \quad (4.4)$$

$s = 1, \dots, m$, where

$$\Delta(w_{r_1}, \dots, w_{r_s}) = \prod_{i < j}^s (w_{r_i} - w_{r_j}); \quad \Delta(w_{r_1}) \equiv 1, \quad (4.5)$$

denotes the Vandermonde determinant. The real constants v_r and w_r , $r = 1, \dots, m+1$, obey the relations

$$\prod_{r=1}^{m+1} v_r = \Delta^{-2}(w_1, \dots, w_{m+1}), \quad \sum_{r=1}^{m+1} w_r = 0. \quad (4.6)$$

In (4.4)

$$C_s = \prod_{s'=1}^m B_{s'}^{-A^{ss'}}, \quad (4.7)$$

where

$$A^{ss'} = \frac{1}{m+1} \min(s, s') [m+1 - \max(s, s')], \quad (4.8)$$

$s, s' = 1, \dots, m$, are components of a matrix inverse to the Cartan one, i.e. $(A^{ss'}) = (A_{ss'})^{-1}$ (see Ch.7 in [43]). Here

$$v_r \neq 0, \quad w_r \neq w_{r'}, \quad r \neq r', \quad (4.9)$$

$r, r' = 1, \dots, m+1$. We note that the solution with $B_s > 0$ may be obtained from the solution with $B_s = 1$ (see [51]) by a certain shift $q^s \mapsto q^s + \delta^s$.

The energy reads [51]

$$E_T = \frac{1}{2} \sum_{s,s'=1}^m A_{ss'} \dot{q}^s \dot{q}^{s'} + \sum_{s=1}^m B_s \exp \left(\sum_{s'=1}^m A_{ss'} q^{s'} \right) = \frac{1}{2} \sum_{r=1}^{m+1} w_r^2. \quad (4.10)$$

If $B_s > 0$, $s \in S$, then all w_r, v_r are real and, moreover, all $v_r > 0$, $r = 1, \dots, m+1$. In a general case $B_s \neq 0$, $s \in S$, relations (4.4)-(4.7) also describe real solutions to eqs. (4.1) for suitably chosen complex parameters v_r and w_r . These parameters are either real or belong to pairs of complex conjugate (non-equal) numbers, i.e., for example, $w_1 = \bar{w}_2$, $v_1 = \bar{v}_2$. When some of B_s are negative, there are also some special (degenerate) solutions to eqs. (4.1) that are not described by relations (4.4)-(4.7), but may be obtained from the latter by certain limits of parameters w_i (see example in the next section).

For the energy (3.39) we get

$$E_{TL} = \frac{1}{2K} E_T = \frac{h}{4} \sum_{r=1}^{m+1} w_r^2. \quad (4.11)$$

Here

$$K_s = K, \quad h_s = h = K^{-1}, \quad (4.12)$$

$s \in S$.

Thus, in the A_m Toda chain case (4.1) eqs. (4.4)-(4.12) should be substituted into relations (3.38) and (3.41)-(3.46) for the general solution.

IV.1 Examples for $d_0 > 2$

Here we consider the case $d_0 > 2$. Let matrix $(h_{\alpha\beta})$ be positively defined and $K = K_s > 0$. Then from the energy constraint (3.38) we get

$$E_{TL} \leq 0 \implies E_T \leq 0. \quad (4.13)$$

In this case

$$c^\alpha = c^i = 0 \iff E_{TL} = E_T = 0, \quad (4.14)$$

$i = 1, \dots, n$, and $\alpha = 1, \dots, l$. When $(h_{\alpha\beta})$ is negatively defined (as it takes place in 12-dimensional theory from [24]) there exist solutions with $E_{TL} > 0$.

IV.1.1 A_1 -case.

Here we consider the case of one “brane”, i.e. $S = \{s\}$. Solving the Liouville equation

$$\ddot{q}^s = -B_s \exp(2q^s), \quad (4.15)$$

with $B_s = \varepsilon_s K_s Q_s^2$, we get

$$f_s(H) = |K_s|^{1/2} |Q_s| \hat{f}_s(H), \quad (4.16)$$

where $H = H(x) > 0$ and

$$\hat{f}_s(H) = \frac{1}{\sqrt{E_T}} \text{ch}(\sqrt{E_T} H), \quad \varepsilon_s K_s > 0, \quad E_T > 0; \quad (4.17)$$

$$\frac{1}{\sqrt{E_T}} \text{sh}(\sqrt{E_T} H), \quad \varepsilon_s K_s < 0, \quad E_T > 0; \quad (4.18)$$

$$\frac{1}{\sqrt{-E_T}} \sin(\sqrt{-E_T} H), \quad \varepsilon_s K_s < 0, \quad E_T < 0; \quad (4.19)$$

$$H, \quad \varepsilon_s K_s < 0, \quad E_T = 0. \quad (4.20)$$

Here

$$E_T = (\dot{q}^s)^2 + B_s \exp(2q^s). \quad (4.21)$$

In a special case $E_T = 0$ this solution agrees with those from [25, 36] if the following redefinition of the harmonic function is performed: $H \mapsto \hat{H}$,

$$\hat{H} = |K_s|^{1/2} |Q_s| H. \quad (4.22)$$

IV.1.2 A_2 -case.

Now we consider the case $m = 2$ (for perfect fluid case see also [52]). Here we put $S = \{1, 2\}$. The solution reads

$$C_1 \exp(-q^1) = v_1 \exp(w_1 u) + v_2 \exp(w_2 u) + v_3 \exp(w_3 u), \quad (4.23)$$

$$\begin{aligned} C_2 \exp(-q^2) &= v_1 v_2 (w_1 - w_2)^2 \exp(-w_3 u) \\ &+ v_2 v_3 (w_2 - w_3)^2 \exp(-w_1 u) + v_3 v_1 (w_3 - w_1)^2 \exp(-w_2 u), \end{aligned} \quad (4.24)$$

where

$$w_1 + w_2 + w_3 = 0, \quad (4.25)$$

$$v_1 v_2 v_3 = (w_1 - w_2)^{-2} (w_2 - w_3)^{-2} (w_3 - w_1)^{-2}. \quad (4.26)$$

and

$$C_1 = (B_1^2 B_2)^{-1/3}, \quad C_2 = (B_2^2 B_1)^{-1/3}. \quad (4.27)$$

Let $K > 0$. Then $E_T \leq 0$ and hence some of B_i should be negative.

Let $B_1 < 0$ and $B_2 < 0$. In the pseudo-Euclidean case, when $\varepsilon[g] = -1$ and all $\theta_a = 1$, this means that $\varepsilon_{s_i} = \varepsilon(I_{s_i}) = -1$, $i = 1, 2$, (see (2.21)), i.e. all p -branes should contain an odd number of time submanifolds.

Let us consider solutions with a negative energy

$$E_T = \frac{1}{2}(w_1^2 + w_2^2 + w_3^2) < 0. \quad (4.28)$$

In this case two of parameters w_i should be complex. Without loss of generality we put

$$w_1 = -2\alpha, \quad w_2 = \alpha + i\beta, \quad w_3 = \alpha - i\beta, \quad (4.29)$$

$$v_2 = v e^{i\theta}, \quad v_3 = v e^{-i\theta}, \quad (4.30)$$

where parameters $\beta \neq 0$, α and $v > 0$ are real and

$$v_1 v^2 = -\frac{1}{4}\beta^{-2}(9\alpha^2 + \beta^2)^{-2}. \quad (4.31)$$

Then relations (4.23) and (4.24) read

$$|C_1| \exp(-q^1) = -v_1 \exp(-2\alpha u) + 2v \exp(\alpha u) \cos(\beta u + \theta), \quad (4.32)$$

$$|C_2| \exp(-q^2) = 4\beta^2 v^2 \exp(2\alpha u) - 2v v_1 (9\alpha^2 + \beta^2) \exp(-\alpha u) \cos(\beta u + \theta + 2\varphi), \quad (4.33)$$

where $|C_1| = (B_1^2 |B_2|)^{-1/3}$, $|C_2| = (B_2^2 |B_1|)^{-1/3}$, $|B_j| = Q_j^2$, $j = 1, 2$, and

$$3\alpha + i\beta = (9\alpha^2 + \beta^2)^{1/2} e^{i\varphi}. \quad (4.34)$$

There exists also a degenerate solution with $E_T = 0$

$$|C_1| \exp(-q^1) = |C_2| \exp(-q^2) = \frac{1}{2}(u - u_0)^2, \quad (4.35)$$

$u_0 = \text{const}$, that may be obtained from the solution (4.32), (4.33) with $\alpha = 0$, $v_1 = -2v$, $v = 1/2\beta^2$, $\theta = -\beta u_0$

$$|C_1| \exp(-q^1) = |C_2| \exp(-q^2) = \frac{2}{\beta^2} \sin^2[\beta(u - u_0)/2], \quad (4.36)$$

in the limit $\beta \rightarrow 0$.

Let us consider an example of the A_2 -solution in $D = 11$ supergravity [44]. We put $n = 3$, $g^3 = -dt \otimes dt$, $d_1 = 2$, $d_2 = 5$, $d_0 = 3$ (metrics g^0, g^1, g^2 are Ricci-flat). The

A_2 -solution, describing a dyon configuration with electric 2-brane and magnetic 5-brane, corresponding to 4-form F and intersecting in 1-dimensional time manifold reads:

$$g = c^{2/9}(\hat{H}^2 g^0 - \hat{H}^{-2} dt \otimes dt) + c^{8/9} g^1 + c^{-4/9} g^2, \quad (4.37)$$

$$F = c\nu_1 d\hat{H}^{-1} \wedge dt \wedge \tau_1 + c\nu_2 (*_0 d\hat{H}) \wedge \tau_1, \quad (4.38)$$

where $\hat{H}$ is the harmonic function on (M_0, g^0) and $\nu_1^2 = \nu_2^2 = 1$. This solution corresponds to the degenerate solution (4.35). Here the following notations $\hat{H} = H(|Q_1||Q_2|)^{1/2}$, $c = (|Q_2|/|Q_1|)^{1/3}$ are adopted. For $c = 1$ this solution coincides with that of [36].

V Cosmological-type solutions

V.1 Solutions with Ricci-flat spaces

Let us consider a “cosmological” case: $d_0 = 1$ and

$$M_0 = \mathbf{R}, \quad g^0 = w du \otimes du, \quad (5.1)$$

where $w = \pm 1$. Since $H(u) = u$ is a harmonic function on (M_0, g^0) we get for the metric and scalar fields from (3.43), (3.45)

$$g = \left(\prod_{s \in S} f_s^{2d(I_s)h_s/(D-2)} \right) \left\{ \exp(2c^0 u + 2\bar{c}^0) w du \otimes du \right. \quad (5.2)$$

$$+ \sum_{i=1}^n \left(\prod_{s \in S} f_s^{-2h_s \delta_{iI_s}} \right) \exp(2c^i u + 2\bar{c}^i) g^i \Big\},$$

$$\exp(\varphi^\alpha) = \left(\prod_{s \in S} f_s^{h_s \chi_s \lambda_{a_s}^\alpha} \right) \exp(c^\alpha u + \bar{c}^\alpha), \quad (5.3)$$

$\alpha = 1, \dots, l$, where $f_s = f_s(u) = \exp(-q^s(u))$ and $q^s(u)$ obey Toda-like equations (3.26).

Relations (3.37) and (3.38) take the form

$$c^0 = \sum_{j=1}^n d_j c^j, \quad \bar{c}^0 = \sum_{j=1}^n d_j \bar{c}^j, \quad (5.4)$$

$$2E = 2E_{TL} + h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^n d_i (c^i)^2 - \left(\sum_{i=1}^n d_i c^i \right)^2 = 0, \quad (5.5)$$

with E_{TL} from (3.39) and all other relations (e.g. constraints (3.29), (3.30)) and relations for forms (3.41) and (3.42) with $H = u$) are unchanged. In a special A_m Toda chain case this solution was considered previously in [42].

V.2 Solutions with one curved space

The cosmological solution with Ricci-flat spaces may be also modified to the following case:

$$\xi_1 \neq 0, \quad \xi_2 = \dots = \xi_n = 0, \quad (5.6)$$

i.e. one space is curved and others are Ricci-flat and

$$1 \notin I_s, \quad \forall s \in S, \quad (5.7)$$

i.e. all “brane” submanifolds do not contain M_1 .

Relation (5.6) modifies the potential (3.17) by adding the term

$$\delta V = \frac{1}{2} w \xi_1 d_1 \exp(2U^1(z)), \quad (5.8)$$

where ($d_1 > 1$)

$$U^1(z) = U_A^1 z^A = -\phi^1 + \gamma_0(\phi), \quad (5.9)$$

$$(U_A^1) = (-\delta_i^1 + d_i, 0). \quad (5.10)$$

For the scalar products we get [31]

$$(U^1, U^1) = \frac{1}{d_1} - 1 < 0, \quad (5.11)$$

$$(U^1, U^s) = 0 \quad (5.12)$$

for all $s \in S$.

The solutions in the case under consideration may be obtained by a little modification of the solution (3.25) (see Appendix)

$$z^A(u) = -\frac{U^{1A}}{(U^1, U^1)} \ln |f_1(u - u_1)| - \sum_{s \in S} \frac{U^{sA}}{(U^s, U^s)} \ln(f_s(u)) + c^A u + \bar{c}^A, \quad (5.13)$$

where

$$f_1(\tau) = R \operatorname{sh}(\sqrt{C_1} \tau), \quad C_1 > 0, \quad \xi_1 w > 0; \quad (5.14)$$

$$R \sin(\sqrt{|C_1|} \tau), \quad C_1 < 0, \quad \xi_1 w > 0; \quad (5.15)$$

$$R \operatorname{ch}(\sqrt{C_1} \tau), \quad C_1 > 0, \quad \xi_1 w < 0; \quad (5.16)$$

$$|\xi_1(d_1 - 1)|^{1/2} \tau, \quad C_1 = 0, \quad \xi_1 w > 0, \quad (5.17)$$

u_1 and C_1 are constants and $R = |\xi_1(d_1 - 1)/C_1|^{1/2}$.

Vectors $c = (c^A)$ and $\bar{c} = (\bar{c}^A)$ satisfy the linear constraints (3.29), (3.30) and also additional constraints

$$U^1(c) = U_A^1 c^A = -c^1 + \sum_{j=1}^n d_j c^j = 0, \quad (5.18)$$

$$U^1(\bar{c}) = U_A^1 \bar{c}^A = -\bar{c}^1 + \sum_{j=1}^n d_j \bar{c}^j = 0. \quad (5.19)$$

The zero-energy constraint reads

$$E = E_1 + E_{TL} + \frac{1}{2} \hat{G}_{AB} c^A c^B = 0, \quad (5.20)$$

where $C_1 = 2E_1(U^1, U^1)$ or, equivalently,

$$C_1 \frac{d_1}{d_1 - 1} = 2E_{TL} + h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=2}^n d_i (c^i)^2 + \frac{1}{d_1 - 1} \left(\sum_{i=2}^n d_i c^i \right)^2. \quad (5.21)$$

From (5.11), (5.13) and

$$U^{1i} = -\frac{\delta_1^i}{d_1}, \quad U^{1\alpha} = 0, \quad (5.22)$$

we get a relation for the metric

$$g = \left(\prod_{s \in S} [f_s(u)]^{2d(I_s)h_s/(D-2)} \right) \left\{ [f_1(u - u_1)]^{2d_1/(1-d_1)} \exp(2c^1 u + 2\bar{c}^1) \right. \quad (5.23)$$

$$\left. \times [w du \otimes du + f_1^2(u - u_1) g^1] + \sum_{i=2}^n \left(\prod_{s \in S} [f_s(u)]^{-2h_s \delta_{1s}} \right) \exp(2c^i u + 2\bar{c}^i) g^i \right\}. \quad (5.24)$$

All other relations are unchanged. (Here $H(u) = u$ and $*_0 dH = w$ in (3.42).) This solution in a special case of A_m Toda chain was obtained earlier in [39].

VI Appendix: Solutions for Toda-like system

Let

$$L = \frac{1}{2} \langle \dot{x}, \dot{x} \rangle - \sum_{s \in S} A_s \exp(2 \langle b_s, x \rangle) \quad (6.1)$$

be a Lagrangian, defined on $V \times V$, where V is n -dimensional vector space over $\mathbf{R}$, $A_s \neq 0$, $s \in S$; $S \neq \emptyset$, and $\langle \cdot, \cdot \rangle$ is non-degenerate real-valued quadratic form on V . Let

$$K_s = \langle b_s, b_s \rangle \neq 0, \quad (6.2)$$

for all $s \in S$.

Then, the Euler-Lagrange equations for the Lagrangian (6.1)

$$\ddot{x} + \sum_{s \in S} 2A_s b_s \exp(2 \langle b_s, x \rangle) = 0, \quad (6.3)$$

have the following solutions

$$x(u) = \sum_{s \in S} \frac{q^s(u) b_s}{\langle b_s, b_s \rangle} + \alpha u + \beta, \quad (6.4)$$

where $\alpha, \beta \in V$,

$$\langle \alpha, b_s \rangle = \langle \beta, b_s \rangle = 0, \quad (6.5)$$

$s \in S$, and functions $q^s(u)$ satisfy the Toda-like equations

$$\ddot{q}^s = -2A_s K_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right), \quad (6.6)$$

with

$$A_{ss'} = \frac{2 \langle b_s, b_{s'} \rangle}{\langle b_{s'}, b_{s'} \rangle}, \quad (6.7)$$

$s, s' \in S$. Let the matrix $(A_{ss'})$ be a non-degenerate one. In this case vectors b_s , $s \in S$, are linearly independent. Then eqs. (6.6) are field equations corresponding to the Lagrangian

$$L_{TL} = \frac{1}{4} \sum_{s, s' \in S} K_s^{-1} A_{ss'} \dot{q}^s \dot{q}^{s'} - \sum_{s \in S} A_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right). \quad (6.8)$$

For the energy corresponding to the solution (6.4) we get

$$E = \frac{1}{2} \langle \dot{x}, \dot{x} \rangle + \sum_{s \in S} \exp(2 \langle b_s, x \rangle) = E_{TL} + \frac{1}{2} \langle \alpha, \alpha \rangle, \quad (6.9)$$

where

$$E_{TL} = \frac{1}{4} \sum_{s, s' \in S} K_s^{-1} A_{ss'} \dot{q}^s \dot{q}^{s'} + \sum_{s \in S} A_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right), \quad (6.10)$$

is the energy function corresponding to the Lagrangian (6.9).

For dual vectors $u^s \in V^*$ defined as $u^s(x) = \langle b_s, x \rangle$, $\forall x \in V$, we have $\langle u^s, u^l \rangle_* = \langle b_s, b_l \rangle$, where $\langle \cdot, \cdot \rangle_*$ is dual form on V^* . The orthogonality conditions (6.5) read

$$u^s(\alpha) = u^s(\beta) = 0, \quad (6.11)$$

$s \in S$.

VII Conclusions

Here we obtained a family of solutions in multidimensional gravity with p -branes generalizing Majumdar-Papapetrou type solutions from [25, 36] in a special case of one harmonic function. These solutions are related to Toda-like systems (of general type) and are defined up to the solutions of Laplace and Toda-type equations. We considered special solutions corresponding to A_m Toda lattices (written in a parametrization of A. Andersen [51]). The general solutions may be also used for other open Toda lattices, e.g. corresponding to B_m , C_m , D_m series. The solutions also contain a class of ‘‘cosmological’’ and spherically symmetric solutions and may be used for investigation of possible black hole and wormhole configurations.

Acknowledgments

This work was supported in part by the Russian Ministry for Science and Technology, Russian Foundation for Basic Research, project SEE, Korea Research Foundation made in the program year of 1997, KOSEF No. 95-0702-04-01-3 and project SEE. V.D.I also thanks the members of Department of Science Education and Basic Science Research Institute of Ewha Womans University (Seoul, Korea) for the kind hospitality during his visit in April-May of 1999.

References

- [1] E. Witten, *Nucl. Phys.* **B 443**, 85 (1995); hep-th/9503124;
P. Townsend, *Phys. Lett.* **B 350**, 184 (1995); hep-th/9612121;
C. Hull and P. Townsend, *Nucl. Phys.* **B 438**, 109 (1995); hep-th/9610167;
P. Horava and E. Witten, *Nucl. Phys.* **B 460**, 506 (1996); hep-th/9510209.
- [2] J.M. Schwarz, Lectures on Superstring and M-theory Dualities, hep-th/9607201;
M.J. Duff, M-theory (the Theory Formerly Known as Strings), hep-th/9608117.
- [3] M.B. Green, J.H. Schwarz and E. Witten, *Superstring Theory*, vol. 1, 2, Cambridge, 1987.
- [4] K.S. Stelle, Lectures on Supergravity p-branes, hep-th/9701088.
- [5] M.J. Duff, R.R. Khuri and J.X. Lu, *Phys. Rep.* **259**, 213 (1995).
- [6] A. Strominger, *Phys. Lett.* **B 383**, 44 (1996); hep-th/9512059.
- [7] M. Cvetič and A.A. Tseytlin, *Nucl. Phys.* **B 478**, 181 (1996).
- [8] A.A. Tseytlin, *Nucl. Phys.* **B 487**, 141 (1997); hep-th/9609212.
- [9] G. Papadopoulos and P.K. Townsend, *Phys. Lett.* **B 380**, 273 (1996); hep-th/9603087.
- [10] A.A. Tseytlin, *Nucl. Phys.* **B 475**, 149 (1996); hep-th/9604035.
- [11] J.P. Gauntlett, D.A. Kastor, and J. Traschen, *Nucl. Phys.* **B 478**, 544 (1996); hep-th/9604179.
- [12] H. Lü, C.N. Pope, and K.W. Xu, Liouville and Toda Solitons in M-theory, hep-th/9604058.
- [13] H. Lü, S. Mukherji, C.N. Pope and K.-W. Xu, Cosmological Solutions in String Theories, hep-th/9610107.
- [14] A. Volovich, *Nucl. Phys.* **B 487** (11), 141 (1997); hep-th/9608095.
- [15] I.Ya. Aref'eva and A.I. Volovich, *Class. Quantum Grav.* **B 14**, 29901 (1997); hep-th/9611026.
- [16] V.D. Ivashchuk and V.N. Melnikov, Intersecting p-brane Solutions in Multidimensional Gravity and M-theory, hep-th/9612089; *Grav. and Cosmol.* **2**, No 4, 204 (1996).
- [17] V.D. Ivashchuk and V.N. Melnikov, *Phys. Lett.* **B 403**, 23 (1997).
- [18] E. Bergshoeff, M. de Roo, E. Eyras, B. Janssen and J.P. van der Schaar, *Class. Quantum Grav.* **14**, 2757 (1997); hep-th/9612095.
- [19] I.Ya. Aref'eva and O.A. Rytchikov, Incidence Matrix Description of Intersecting p-brane Solutions, *Preprint* SMI-25-96, hep-th/9612236.
- [20] R. Argurio, F. Englert and L. Hourant, *Phys. Lett.* **B 398**, 2991 (1997); hep-th/9701042.
- [21] I.Ya. Aref'eva, M.G. Ivanov and O.A. Rytchikov, Properties of Intersecting p-branes in Various Dimensions, *Preprint* SMI-05-97, hep-th/9702077.

- [22] I.Ya. Aref'eva, M.G. Ivanov and I.V. Volovich, Non-Extremal Intersecting p-Branes in Various Dimensions, hep-th/9702079; *Phys. Lett. B* **406**, 44 (1997).
- [23] N. Ohta, Intersection Rules for Non-extreme p-branes, hep-th/9702164.
- [24] N. Khvengia, Z. Khvengia, H. Lü and C.N. Pope, *Class. Quantum Grav.* **15**, 759 (1998); hep-th/9703012.
- [25] V.D. Ivashchuk and V.N. Melnikov, Sigma-model for the Generalized Composite p-branes, hep-th/9705036; *Class. Quantum Grav.* **14**, 3001 (1997); Corrigenda *ibid.* **15** (12), 3941 (1998).
- [26] V.D. Ivashchuk, V.N. Melnikov and M. Rainer, Multidimensional σ -models with Composite Electric p -branes, gr-qc/9705005; *Grav. and Cosmol.* **4**, No 1 (13), (1998).
- [27] K.A. Bronnikov, M.A. Grebeniuk, V.D. Ivashchuk and V.N. Melnikov, Integrable Multi-dimensional Cosmology for Intersecting p -branes, *Grav. and Cosmol.* **3**, No 2(10), 105 (1997).
- [28] H. Lü, J. Maharana, S. Mukherji and C.N. Pope, Cosmological Solutions, p-branes and the Wheeler De Witt Equation, hep-th/9707182.
- [29] M.A. Grebeniuk, V.D. Ivashchuk and V.N. Melnikov, Integrable Multidimensional Quantum Cosmology for Intersecting p-Branes, *Grav. and Cosmol.* **3**, No 3 (11), 243 (1997), gr-qc/9708031.
- [30] K.A. Bronnikov, U. Kasper and M. Rainer, Intersecting Electric and Magnetic p -Branes: Spherically Symmetric Solutions, gr-qc/9708058.
- [31] V.D. Ivashchuk and V.N. Melnikov, Multidimensional Classical and Quantum Cosmology with Intersecting p -branes, *J. Math. Phys.*, **39**, 2866 (1998); hep-th/9708157,
- [32] D. Youm, submitted to *Phys. Rep*, hep-th/9710046.
- [33] K.A. Bronnikov, V.D. Ivashchuk and V.N. Melnikov, The Reissner-Nordström Problem for Intersecting Electric and Magnetic p -branes, gr-qc/9710054; *Grav. and Cosmol.*, **3**, No 3(11), 203 (1997).
- [34] K.A. Bronnikov, Block-orthogonal Brane systems, Black Holes and Wormholes, hep-th/9710207; *Grav. and Cosmol.* **4**, No 1 (13), 49 (1998).
- [35] D.V. Gal'tsov and O.A. Rytchkov, Generating Branes via Sigma models, hep-th/9801180.
- [36] V.D. Ivashchuk and V.N. Melnikov, Madjumdar-Papapetrou Type Solutions in Sigma-model and Intersecting p-branes, *Class. Quantum Grav.* **16**, 849 (1999); hep-th/9802121.
- [37] V.D. Ivashchuk, S.-W. Kim and V.N. Melnikov, Hyperbolic Kac-Moody Algebra from Intersecting p -branes, to appear in *J. Math. Phys.*; hep-th/9803006.
- [38] M.A. Grebeniuk and V.D. Ivashchuk, Sigma-model Solutions and Intersecting p-branes Related to Lie Algebras, *Phys. Lett. B* **442**, 125 (1998); hep-th/9805113.

- [39] V.R. Gavrilov and V.N. Melnikov, Toda Chains with Type A_m Lie Algebra for Multidimensional Classical Cosmology with Intersecting p -branes, In : Proceedings of the International seminar "Current topics in mathematical cosmology", (Potsdam, Germany , 30 March - 4 April 1998), Eds. M. Rainer and H.-J. Schmidt, World Scientific, 1998, p. 310; hep-th/9807004.
- [40] V.D. Ivashchuk and V.N. Melnikov, Multidimensional Cosmological and Spherically Symmetric Solutions with Intersecting p -branes, gr-qc/9901001;
Cosmological and Spherically Symmetric Solutions with Intersecting p -branes, to appear in J. Math. Phys.
- [41] S. Cotsakis, V.D. Ivashchuk and V.N. Melnikov, P-branes Black Holes and Post-Newtonian Approximation, *Grav. and Cosmol.* **5**, No 1 (17), (1999); gr-qc/9902148.
- [42] V.R. Gavrilov and V.N. Melnikov, Toda Chains Associated with Lie Algebras A_m in Multidimensional Gravitation and Cosmology with Intersecting p -branes, submitted to Theor. Math. Phys. (in Russian).
- [43] J. Fuchs and C. Schweigert, Symmetries, Lie algebras and Representations. A graduate course for physicists (Cambridge University Press, Cambridge, 1997).
- [44] E. Cremmer, B. Julia, J. Scherk. *Phys. Lett.* **B 76**, 409 (1978).
- [45] S.D. Majumdar, *Phys. Rev.* **72**, 930 (1947);
A. Papapetrou, *Proc. R. Irish Acad.* **A51**, 191 (1947).
- [46] G. Neugebauer and D. Kramer, *Ann. der Physik (Leipzig)* **24**, 62 (1969).
- [47] D. Kramer, H. Stephani, M. MacCallum, and E. Herlt, *Exact Solutions of the Einstein Field Equations*, CUP, 1980.
- [48] G. Clément, *Gen. Rel. and Grav.* **18**, 861 (1986); *Phys. Lett.* **A 118**, 11 (1986).
- [49] V.D. Ivashchuk, V.N. Melnikov and A.I. Zhuk, *Nuovo Cimento* **B 104**, 575 (1989).
- [50] M. Toda, *Progr. Theor. Phys.* **45**, 174 (1970); *Theory of Nonlinear Lattices* (Springer-Verlag, Berlin, 1981).
- [51] A. Anderson, *J. Math. Phys.* **37**, 1349 (1996); hep-th/9507092.
- [52] V.R. Gavrilov, V.D. Ivashchuk and V.N. Melnikov, *J. Math. Phys.* **36**, 5829 (1995).